

Cyclic universe due to phantom and quintessence

Changjun Gao*

*The National Astronomical Observatories, Chinese Academy of Sciences, Beijing, 100012, China and
State Key Laboratory of Theoretical Physics, Institute of Theoretical Physics,
Chinese Academy of Sciences, Beijing 100190, China*

Youjun Lu†

The National Astronomical Observatories, Chinese Academy of Sciences, Beijing, 100012, China

You-Gen Shen‡

Shanghai Astronomical Observatory, Chinese Academy of Sciences, Shanghai 200030, China

(Dated: August 15, 2014)

We explore a cyclic universe due to phantom and quintessence fields. We find that, in every cycle of the evolution of the universe, the phantom dominates the cosmic early history and quintessence dominates the cosmic far future. In this model of universe, there are infinite cycles of expansion and contraction. Different from the inflationary universe, the corresponding cosmic space-time is geodesically complete and quantum stable. But similar to the Cyclic Model, the flatness problem, the horizon problem and the large scale structure of the universe can be explained in this cyclic universe.

PACS numbers: 98.80.Jk, 04.40.Nr, 04.50.+h, 11.25.Mj

I. MOTIVATION

The theory of inflationary cosmology has become the standard paradigm for the early universe, which was firmly established in the past several decades. Inflationary cosmology not only naturally solves the flatness problem, the horizon problem and the magnetic monopole problem, but also elegantly explains the large scale structure of the universe. Despite of its success, it may be not perfect yet and suffer from some conceptual problems as follows.

First, a few common questions may be almost always asked as follows: What is the inflaton field? How to fine-tune the scalar potential in order that it is sufficiently flat? How did the universe begin and why did it start to inflate? These questions are of great importance for the theory consummation. Second, inflationary cosmology suffers from the trans-Planckian problem [1]. In inflationary cosmology, the fluctuations observed in the cosmic microwave background had wavelengths at the beginning of inflation smaller than the Planck scale. It is generally believed that the quantum gravity would become important at sub-Planckian scales. So the quantum gravity may produce different distribution on sub-Planckian wavelengths. This different distribution would be inflated and generate an uncertain correction to the predictions for the cosmic microwave background anisotropy. Third, Borde et al [2] showed that the inflationary space-time is geometrically incomplete in the past directions of the

history of the universe as the expansion rate averaged along the geodesic is positive, i.e., $\langle H \rangle > 0$. This incompleteness theorem relies on neither the energy condition nor the Einstein's equation. In order to solve these problems, a number of alternative scenarios, e.g., the cyclic universe [3, 4], the bounce universe [5] and the emergent universe [6–11] are proposed.

In the Cyclic Model [3], the universe undergoes an endless sequence of cosmic expansion and contraction. The cosmic temperature and energy density are all finite at each transition point from contraction to expansion or from expansion to contraction. The Cyclic Model can solve not only the flatness problem, the horizon problem, the large scale structure problem, but also the Big-Bang singularity problem, the current cosmic accelerating problem and so on. Therefore, the cyclic model may present a complete description of all phases of the cosmic evolution.

The cyclic universe model [3, 4] is originally constructed in the brane-world scenario. In this paper, we alternatively construct a cyclic universe by simply introducing a phantom matter and quintessence matter. In this cyclic universe, there are infinite cycles of expansion and contraction. Different from the inflationary universe, the corresponding cosmic space-time is geometrically complete and quantum stable. Similar to the Cyclic Model, the flatness problem, the horizon problem and the large scale structure of the universe can be explained in this cyclic universe.

This paper is organized as follows. In section II, we present the energy density for the phantom and quintessence, respectively. Then, by taking into account both the relativistic matter and dust matter, we construct a cyclic model for the universe and investigate its cyclic evolution in section III. The total cosmic en-

*Electronic address: gaojcj@bao.ac.cn

†Electronic address: luyj@bao.ac.cn

‡Electronic address: ygshen@center.shao.ac.cn

ergy density is zero at the transition points, either from expansion to contraction or from contraction to expansion. Therefore, the Big-bang and Big-crunch singularity problems are automatically vanishing. In section IV, we demonstrate that the cosmic space-time is geodesically complete for both time-like and null geodesics. The particle and the event horizon are all infinite in this space-time. So the horizon problem is solved. In section V, we discuss the variation of entropy of the cyclic universe and the violation of second law of thermodynamics due to the presence of phantom matter. In section VI, using the Wheeler-DeWitt equation, we find that the system is quantum stable although a negative scalar potential is present. In section VII, we verify that the primordial scale-invariant power spectrum problem could be generated in this cyclic model. Finally, conclusions and discussions are given in section VIII. Throughout this paper, we adopt the system of units in which $G = c = \hbar = 1$ and the metric signature $(-, +, +, +)$.

II. PHANTOM AND QUNITESSENCE

Phantom matter is introduced into in the study of cosmic evolution by Caldwell in Ref. [12]. The Lagrangian of phantom matter takes the form

$$\mathcal{L}_\phi = -\frac{1}{2}\nabla_\mu\phi\nabla^\mu\phi - V(\phi), \quad (1)$$

with ϕ the phantom field and $V(\phi)$ the phantom potential. For our purpose, we shall simply take a vanishing potential. Then it is straightforward to show the phantom ϕ with the Lagrangian

$$\mathcal{L}_\phi = -\frac{1}{2}\nabla_\mu\phi\nabla^\mu\phi, \quad (2)$$

contributes the energy density

$$\rho_\phi = -\frac{\rho_{\phi_0}}{a^6}, \quad (3)$$

in the background of spatially flat Friedmann-Robertson-Walker Universe. Here ρ_{ϕ_0} is an integration constant which has the physical meaning of the minus of energy density when the cosmic scale factor $a = 1$.

Compared to the Lagrangian of quintessence field [13], the sign before the kinetic term of phantom is not positive, but negative. So the kinetic energy density of phantom becomes negative. For simplicity, we shall consider the quintessence field with the Lagrangian as follows

$$\mathcal{L}_\psi = \frac{1}{2}\nabla_\mu\psi\nabla^\mu\psi + \frac{\rho_\Lambda}{3} [1 - 2\sin(4\sqrt{\pi}\psi)], \quad (4)$$

which contributes the energy density

$$\rho_\psi = \rho_\Lambda - \rho_{\psi_0}a^2. \quad (5)$$

Here ρ_Λ and ρ_{ψ_0} are two integration constants. The reason for these specific choices will be given in the next section and we shall show that, with these terms, a cyclic universe is constructed.

III. CYCLIC UNIVERSE

Now the Friedmann equation could be written as follows

$$3H^2 = 8\pi \left(-\frac{\rho_{\phi_0}}{a^6} + \frac{\rho_{r_0}}{a^4} + \frac{\rho_{m_0}}{a^3} + \frac{\rho_k}{a^2} + \rho_\Lambda - \rho_{\psi_0}a^2 \right). \quad (6)$$

On the right hand side of Eq. (6), the energy densities are from phantom field, relativistic matter, dust matter (which includes dark matter and baryonic matter), cosmic spatial curvature and quintessence, respectively. The parameters ρ_{ϕ_0} , ρ_{r_0} , ρ_{m_0} , ρ_k , ρ_Λ , and ρ_{ψ_0} represent the present-day energy density of each component, respectively. The parameter ρ_k can be positive, negative or 0, and the other five parameters are all positive.

Now we could understand why we introduce the negative scalar potential. According to the Friedmann equation for the standard Λ CDM cosmology

$$3H^2 = 8\pi \left(\frac{\rho_{r_0}}{a^4} + \frac{\rho_{m_0}}{a^3} + \frac{\rho_k}{a^2} + \rho_\Lambda \right), \quad (7)$$

the evolution of the scale factor a is controlled by the relativistic matter if a is sufficiently small, and $a \propto \sqrt{t}$. At the Big-bang singularity, $t = 0$, the scale factor vanishes and the energy density is divergent. In order to erase the Big-bang singularity and turn the universe from contraction to expansion, namely, making the velocity of the universe $\dot{a}$ varying from negative (contracting universe) to positive (expanding universe), it is necessary to introduce the negative energy density. At the turning point, we should have $\dot{a} = 0$.

For simplicity, the negative energy density may be assumed to scale as a^{-n} with $n > 4$ such that the velocity of the universe $\dot{a}$ could cross zero at the minimum of the scale factor. We take $n = 6$ in this paper. Similarly, with the increasing of the scale factor, dark energy dominates finally. In order to let the universe contract at sometime in the future, it is also necessary to introduce negative energy density that dominates at large a , and this density may be assumed to scale as a^m with $m > 0$ such that the velocity of the universe $\dot{a}$ could cross zero at the maximum of the scale factor. In this study, we simply take $m = 2$.

Given the Friedmann equation Eq. (6), and the equation of conservation for energy, we derive the acceleration equation

$$6\frac{\ddot{a}}{a} = -8\pi \left(-4\frac{\rho_{\phi_0}}{a^6} + 2\frac{\rho_{r_0}}{a^4} + \frac{\rho_{m_0}}{a^3} - 2\rho_\Lambda + 4\rho_{\psi_0}a^2 \right). \quad (8)$$

For the present-day universe, we have the Friedmann equation

$$3H_0^2 = 8\pi\rho_0, \quad (9)$$

where H_0 and ρ_0 are the present-day Hubble parameter and total cosmic energy density.

Dividing the acceleration equation Eq. (8) by Eq. (9), we obtain

$$\frac{2}{H_0^2} \cdot \frac{\ddot{a}}{a} = \frac{4\Omega_{\phi_0}}{a^6} - \frac{2\Omega_{r_0}}{a^4} - \frac{\Omega_{m_0}}{a^3} + 2\Omega_{\Lambda} - 4\Omega_{\psi_0}a^2, \quad (10)$$

where the dimensionless constants (the ratio of each component in the present-day cosmic energy density) are defined as

$$\begin{aligned} \Omega_{\phi_0} &\equiv \frac{\rho_{\phi_0}}{\rho_0}, & \Omega_{r_0} &\equiv \frac{\rho_{r_0}}{\rho_0}, & \Omega_{m_0} &\equiv \frac{\rho_{m_0}}{\rho_0}, \\ \Omega_{\Lambda} &\equiv \frac{\rho_{\Lambda}}{\rho_0}, & \Omega_{\psi_0} &\equiv \frac{\rho_{\psi_0}}{\rho_0}, & \Omega_k &\equiv \frac{\rho_k}{\rho_0}. \end{aligned} \quad (11)$$

We can let the cosmic time t absorb H_0 in Eq. (10). Then the unit of cosmic time is the present-day Hubble age

$$\tau = \frac{1}{H_0}. \quad (12)$$

Eq.(10) is reduced to

$$\frac{2\ddot{a}}{a} = \frac{4\Omega_{\phi_0}}{a^6} - \frac{2\Omega_{r_0}}{a^4} - \frac{\Omega_{m_0}}{a^3} + 2\Omega_{\Lambda} - 4\Omega_{\psi_0}a^2. \quad (13)$$




FIG. 1: Evolution of the scale factor $\ln a$ with the cosmic time (in unit of the present-day Hubble age).

As an example, we consider the cosmology model with the following values of parameters, $\Omega_{r_0} = 8.1 \cdot 10^{-5}$, $\Omega_{m_0} = 0.27$, $\Omega_{\Lambda} = 0.73$, and $\Omega_k = 0$, which are consistent with current observations [14]. As for the values of Ω_{ϕ_0} and Ω_{ψ_0} , they must be sufficiently small in order to be not contradict with the well constrained standard Λ CDM model. However, with decreasing scale factor, the universe is finally dominated by both the Ω_{ϕ_0} term and the radiation. A maximum energy density appears in this epoch. It is generally believed that the Planck energy density,

$$\Omega_{\text{Planck}} \simeq 10^{123}, \quad (14)$$




FIG. 2: Velocity $v \equiv 10^{-4}\dot{a}$ of the universe with the cosmic time t (in unit of the present-day Hubble age).




FIG. 3: Evolution of the total cosmic energy density (ρ) with the cosmic time t . The unit of t and ρ is the present-day Hubble age and total cosmic energy density, respectively.

is the highest one. So Ω_{ϕ_0} is constrained to be in the order of $\Omega_{\phi_0} \simeq 10^{-68}$. For Ω_{ψ_0} , we simply set $\Omega_{\psi_0} \simeq 10^{-10}$, which is not inconsistent with astronomical observations.

Figures 1 and 2 show the evolution of the scale factor $\ln a$ and the velocity $v \equiv 10^{-4}\dot{a}$ of the universe with respect to the cosmic time t . Apparently there are infinite cycles, expansion and contraction, in the evolution of the universe. The present-day universe locates at $\ln a = 0$ as we have set $a = 1$ for the present-day universe. The minimum and the maximum of the scale factor are $a_{\min} \simeq (\Omega_{\phi_0}/\Omega_{r_0})^{1/2} \sim e^{-73}$ and $a_{\max} \simeq (\Omega_{\psi_0}/\Omega_{\Lambda})^{1/2} \sim e^{11}$, respectively (see Fig. 1). Note here that what characters the physical size of the universe is not the scale factor, but

the apparent horizon, event horizon and particle horizon. We shall come back to this point in the next section.

Fig. 3 shows the evolution of total cosmic energy density ρ with respect to the cosmic time. The maximum and minimum of the energy density are the Planck density and zero, respectively. As seen from Figure 3, the universe contracts from B to C and expands from C to D with the increasing of time. The contracting and expanding phase constitute a complete cycle of evolution. In order to understand the evolution in detail, let's start from point A , which represents the time one cycle period before the present day (point Now in Fig. 3). In the first several Hubble ages (a plateau), the universe is dominated by the term Ω_Λ . Then the term $-\Omega_{\psi_0}a^2$ grows and the total cosmic energy density evolves as $\rho \simeq \Omega_\Lambda - \Omega_{\psi_0}a^2$. The energy density decreases with the expansion of the universe. When the density vanishes, we find from the Friedmann equation Eq. (6) that the velocity $\dot{a}$ of the universe also vanishes but the acceleration of the universe $\ddot{a}$ becomes negative (from Eq. (13)). So after crossing point B , the universe evolves into contracting phase. B (and D) corresponds to the maximum of the scale factor as clearly shown in Fig. 1.

In the contracting phase (from B to C), the universe is first dominated by $\Omega_\Lambda - \Omega_{\psi_0}a^2$, and then by Ω_Λ (the plateau), by the matter, by the radiation, and by radiation and term Ω_{ϕ_0} in a sequence. The maximum of energy density appears at $a \simeq \frac{\sqrt{6}}{2}a_{\min}$. With the increasing of term Ω_{ϕ_0} , the comic density decreases sharply to 0 at $a = a_{\min}$. At the minimum of the scale factor, both the cosmic density and the cosmic velocity vanishes (point C). After point C , the universe goes into the expanding phase. At the turning point C , we have $\dot{a} = 0$ and $\ddot{a} > 0$ which follows from Eq. 6 and Eq. 13.

In the expanding phase (from C to D), the universe is firstly dominated by both the chameleon scalar and the radiation, $\rho \simeq \Omega_{r_0}/a^4 - \Omega_{\phi_0}/a^6$. Since the density of chameleon scalar decreases quickly with the expansion of the universe, the cosmic energy density grows significantly. Once the cosmic energy density crosses the maximum, the universe evolves as the standard Λ CDM model (before arriving at point D). The present-day universe locates at $\rho = 1$ (one unit of present-day cosmic energy density, ρ_0). It is apparent the age of the universe (from the maximum energy density to Now) is within one Hubble age which is consistent with observations.

IV. GEODESIC COMPLETE SPACE-TIME AND HORIZONS

A. geodesic complete space-time

In this sub-section, we show that the space-time of the cyclic universe constructed in section III is complete for both null and time-like geodesics. Therefore, there is no singularity in this space-time. To this end, let's first derive the equation of motion for a photon and a

massive particle. The metric of spatially-flat Friedmann-Robertson-Walker Universe is give by

$$ds^2 = dt^2 - a(t)^2 (dx^2 + dy^2 + dz^2) . \quad (15)$$

Then the Lagrangian for a photon and a massive particle propagating in this space-time is given by [15]

$$\mathcal{L} = \frac{1}{2} [\dot{t}^2 - a^2 (\dot{x}^2 + \dot{y}^2 + \dot{z}^2)] \equiv \epsilon , \quad (16)$$

where dot denotes the derivative with respect to the affine parameter τ , and τ is exactly the proper time of a massive particle. We have $\epsilon = 0$ and $\epsilon = 1/2$ for the photon and massive particle (with mass of unit one), respectively [15]. Using the Euler-Lagrange equation, we obtain the equation of geodesics

$$\dot{t}^2 - \frac{\zeta^2}{a^2} - 2\epsilon = 0 , \quad (17)$$

where $\epsilon = 0$ for the null geodesics and $\epsilon = 1/2$ for the time-like geodesics, and ζ is an integration constant that represents the momentum of the particle. Therefore, τ is integrated as

$$\tau = + \int_0^{+\infty} \frac{1}{\sqrt{2\epsilon + \frac{\zeta^2}{a^2}}} dt = \infty , \quad (18)$$

for the future-pointing geodesics, and

$$\tau = - \int_0^{-\infty} \frac{1}{\sqrt{2\epsilon + \frac{\zeta^2}{a^2}}} dt = \infty , \quad (19)$$

for the past-pointing geodesics, respectively. Equations (18) and (19) show that the space-time is complete for both null and time-like geodesics.

B. horizons

In general, there are three horizons in the Friedmann-Robertson-Walker space-time, i.e., the particle horizon, event horizon and apparent horizon. In a universe with a finite age, a photon would only propagate a finite distance in that finite time and the volume of space from which we can receive information at a given time is also limited. The boundary of this volume is called the particle horizon which is given by

$$d_p = a \int_{t_i}^t \frac{dt}{a} , \quad (20)$$

where t_i is the beginning of the universe. For the cyclic universe presented above, we have $t_i = -\infty$ and $a_{\min} \leq a \leq a_{\max}$. So the corresponding particle horizon is infinite.

The event horizon is defined as the complement of the particle horizon. It encloses the set of points from which

photons sent at a given time t will never be received by an observer in the future. For our cyclic universe, we find the physical size of the event horizon is also infinite,

$$d_e = a \int_t^{+\infty} \frac{dt}{a} = \infty. \quad (21)$$

The apparent horizon is a marginally trapped surface with vanishing expansion and has been argued to be a causal horizon for a dynamical space-time. The apparent horizon is associated with the Hawking temperature, gravitational entropy and other thermodynamical aspects [16–18]. The first law of thermodynamics for the apparent horizon has been derived not only in general relativity but also in several alternative theories of gravity, including the Lovelock, nonlinear, scalar-tensor, and braneworld theories [19–25]. Therefore, it is of great importance for us to consider the apparent horizon in the cyclic model of the universe. For the spatially flat universe, the physical size of the apparent horizon is given by

$$d_a = \frac{1}{H}. \quad (22)$$

We recognize that it is exactly the Hubble scale or the Hubble horizon. For our cyclic universe, we find there is a minimum of the physical size of the apparent horizon, namely, the Planck length. At the turning points from expansion to contraction (and from contraction to expansion), the Hubble radius is infinite. Since the particle horizon and event horizon are infinite, the cosmic horizon problem is solved.

V. ENTROPY OF THE CYCLIC UNIVERSE AND VIOLATION OF SECOND LAW OF THERMODYNAMICS

The cyclic universe constructed above is precisely cyclic in the evolution of the scale factor of the universe. The early work of Tolman [26] in the 1930s showed that such a universe is in fact impossible as long as the usual laws of thermodynamics hold. The second law implies that the entropy always increases, and this implies that from one cycle to the next, the universe must grow. We find that the second law of thermodynamics is actually violated in the presence of phantom matter. So Tolman's proof does not apply. In the next, we shall show this point in detail.

Many researches [27] on the relation between thermodynamics and gravitation theories indicate that the entropy of the universe is proportional to the apparent horizon area of the universe, which is also implied by the holographic principle

$$S = \frac{1}{4}A = \frac{1}{4} \cdot 4\pi d_a^2 = \frac{\pi}{H^2}, \quad (23)$$

where d_a is the radius of cosmic apparent horizon defined by Eq. (22). Then the variation of cosmic entropy with respect to the cosmic time t is

$$\begin{aligned} \frac{dS}{dt} &= -\frac{2\pi}{H^3} \dot{H} \\ &= 3\pi d_a (1 + w'). \end{aligned} \quad (24)$$

Here w' is the equation of state defined as $w' = \frac{\sum_i p_i}{\sum_i \rho_i}$, where i represents the i -th component of matters. In the last step, we have used the Einstein equation, $\dot{H} = -4\pi(\rho + p)$. Eq. (24) reveals that for the dust matter ($w' = 0$), relativistic matter ($w' = 1/3$) and quintessence matter ($-1 \leq w' \leq 1$) dominated expanding universe, the entropy of the universe is always increasing. However, for a phantom dominated expanding universe ($w' < -1$), the entropy of the universe is decreasing. This is very much similar to the finding of Babichev et al [28] that the black hole mass (namely, the black hole entropy, $S \propto M^2$) decreases with the accretion of phantom matter. Actually, Ref. [29] has noted that the entropy of the phantom matter is negative. Then we conclude that the second law of thermodynamics in the phantom dominated universe is violated. This is closely related to the violation of null energy condition ($\rho + p > 0$) by phantom matter.

VI. QUANTUM STABILITY

In this section, we shall consider the quantum stability of this cyclic universe model. In the approximation of minisuperspace, the wave function of the universe depends only on one freedom, the scale factor a . The Hamiltonian of the universe is given by

$$\mathcal{H} = -\frac{1}{3\pi a} [p_a^2 + U(a)], \quad (25)$$

where

$$p_a = -\frac{3\pi}{2} a \dot{a}, \quad (26)$$

is the momentum conjugate to a and the potential $U(a)$ is given by

$$U(a) = \frac{9\pi^2}{4} a^2 \left(1 - \frac{8\pi}{3} a^2 \rho \right), \quad (27)$$

where ρ is total energy density of the universe. Then the Hamiltonian constraint $\mathcal{H} = 0$ gives exactly the Friedmann equation, Eq. (6).

In the theory of quantum cosmology, the universe is described by a wave function $\Psi(a)$. The conjugate momentum p_a becomes a differential operator $-i\frac{d}{da}$ and the Hamiltonian constraint is replaced by the Wheeler-DeWitt (WDW) equation [30]

$$\mathcal{H}\Psi = 0, \quad (28)$$

i.e.,

$$\left(-\frac{d^2}{da^2} + U \right) \Psi = 0. \quad (29)$$




FIG. 4: The evolution of the potential $U(a)$. Points C and D mark the minimum and maximum of the scale factor a , respectively. The region between C and D is classically allowed. As the potential is divergent when $a \rightarrow 0$ and $a \rightarrow \infty$, the universe can not tunnel to the classically forbidden regions.

Fig. 4 shows the potential U with respect to the scale factor a . There is an oscillating region between the classical turning points C and D which correspond to the minimum and maximum of the scale factor, respectively. The region between C and D is classically allowed.

The semiclassical tunnelling probability as the universe bounces at C can be determined from

$$\mathcal{P} \sim e^{-2S_{WKB}}, \quad (30)$$

where the tunnelling action are

$$S_{WKB} = \int_0^{a_{min}} \sqrt{U} da = +\infty, \quad (31)$$

Since the tunnelling probability is vanishing, the cyclic universe is able to last forever. In other words, this model of cyclic universe is not immune to collapse and it is indeed quantum stable.

VII. GENERATION OF SCALE-INVARIANT POWER SPECTRUM

In the Cyclic Model [3] of the universe, because the equation of state $w' \gg 1$ during the contraction phase, the universe is homogeneous, isotropic and flat [33] with a scale-invariant spectrum of density perturbations [31, 32]. The $w' \gg 1$ condition also ensures that anisotropy is small and first order perturbation theory remains valid until just before the bounce [33, 34].

On the other hand, it is well-known that in order to generate scale invariant curvature perturbations in a contracting universe, two scalar fields are necessary. For the most recent models, see Ref. [35]. Fig. 5 shows the evolution of w' with the natural logarithm of the scale factor ($\ln a$). As shown in Fig. 5, $w' \gg 1$ in the vicinity of bouncing point B (or D) (see Fig. (1), Fig. (2) or Fig. (3)). Thus we conclude that the flatness of spacetime and the scale-invariant primordial power spectrum can also be generated in the present cyclic universe model. Furthermore, we find the scale-invariant power spectrum could be generated in the vicinity of bouncing point C (or E). In this section, we shall study the generation of scale-invariant power spectrum in the vicinity of bouncing points.




FIG. 5: Evolution of the equation of state w' with the natural logarithm of scale factor $\ln a$. We have $|w'| \gg 1$ in the vicinity of the turning point from expansion to contraction (and from contraction to expansion)

A. Bouncing point C (or E)

Around the bouncing point C (or E, see Fig. (1), Fig. (2) or Fig. (3)), the evolution of the universe is governed by the following Friedmann equation

$$3H^2 = 8\pi \left(-\frac{\rho_{\phi_0}}{a^6} + \frac{\rho_{r_0}}{a^4} \right). \quad (32)$$

Rewrite above equation of motion with the conformal time η and rescale both the cosmic comoving time and the scale factor, we derive the evolution of the scale factor

$$a = \sqrt{1 + \eta^2}. \quad (33)$$

Here the minimum of the scale factor is rescaled to be unit one and $\eta = 0$ denotes the bouncing moment. In the vicinity of bouncing point, namely, $|\eta| \ll 1$, the comoving Hubble parameter $\mathcal{H}$ could be safely approximate as

$$\mathcal{H} = \frac{\eta}{1 + \eta^2} \simeq \eta. \quad (34)$$

Fig. (5) tells us the total equation of state in this epoch is very much smaller than -1 . So we can effectively use the phantom matter to describe the matter content of the universe. Because the evolution of the linear perturbations is determined by the background evolution, namely, the evolution of the Hubble parameter, we need not to know the expression of phantom potential in great detail.

The equation of motion for the adiabatic scalar perturbations is derived in Ref. [36]

$$v_k'' + \left(k^2 - \frac{z''}{z} \right) v_k = 0, \quad (35)$$

where

$$z \equiv \frac{a}{H} \sqrt{|\rho + p|}, \quad (36)$$

and the prime denotes the derivative with respect to the cosmic comoving time. Taking into account the Friedmann equation, the acceleration equation and the expression of Hubble parameter, Eq (34), we have

$$\begin{aligned} z &\equiv \frac{a}{H} \sqrt{|\rho + p|} \propto \frac{a}{H} \sqrt{|\dot{H}|} \\ &= \frac{a \left[\frac{1}{a} \left(\frac{\mathcal{H}}{a} \right)' \right]^{1/2}}{\frac{\mathcal{H}}{a}} \simeq \frac{1}{\eta}. \end{aligned} \quad (37)$$

Here dot denotes the derivative with respect to cosmic time t and $\mathcal{H}$ is the Hubble parameter in terms of the comoving time η . So

$$\frac{z''}{z} = \frac{2}{\eta^2}. \quad (38)$$

Then the equation of motion for the adiabatic scalar perturbations turns out to be

$$v_k'' + \left(k^2 - \frac{2}{\eta^2} \right) v_k = 0. \quad (39)$$

Refs. [37] and [38] have shown that above equation lead to exact scale invariant power spectrum.

B. Bouncing point B (or D)

Similarly, around the bouncing point **B** (or **D**), the Friedmann equation is given by

$$3H^2 = 8\pi (\rho_\Lambda - \rho_{\psi_0} a^2). \quad (40)$$

Rescale the cosmic comoving time and the scale factor, we obtain the evolution of the scale factor

$$a = \frac{1}{\sqrt{1 + \eta^2}}. \quad (41)$$

Here the maximum of the scale factor is rescaled to be unit one and $\eta = 0$ denotes the bouncing moment. So

in the vicinity of bouncing point $|\eta| \ll 1$, the comoving Hubble parameter $\mathcal{H}$ is

$$\mathcal{H} = -\frac{\eta}{1 + \eta^2} \simeq -\eta. \quad (42)$$

The total equation of state in this epoch is very much larger than unit one. Therefore we can effectively use the quintessence matter (but negative scalar potential) to describe the matter content of the universe while irrespective of the knowledge of the quintessence potential.

We have

$$\begin{aligned} z &\equiv \frac{a}{H} \sqrt{|\rho + p|} \propto \frac{a}{H} \sqrt{|\dot{H}|} \\ &= \frac{a \left[\frac{1}{a} \left(\frac{\mathcal{H}}{a} \right)' \right]^{1/2}}{\frac{\mathcal{H}}{a}} \simeq \frac{1}{\eta}. \end{aligned} \quad (43)$$

and

$$\frac{z''}{z} = \frac{2}{\eta^2}. \quad (44)$$

Then the equation of motion for the adiabatic scalar perturbations is also give by

$$v_k'' + \left(k^2 - \frac{2}{\eta^2} \right) v_k = 0. \quad (45)$$

In other words, the scale invariant power spectrum could be also produced in the vicinity of bouncing point **B** (or **D**).

VIII. CONCLUSION AND DISCUSSION

In conclusion, we have explored a cyclic model for the universe by simply considering a phantom field and a quintessence field. In every cycle of the cosmic evolution, the phantom field is dominant in the very early universe and the quintessence field is dominant in the far future.

There are infinite cycles of cosmic expansion and contraction in this model. At the turning points from expansion to contraction or from contraction to expansion, the cosmic energy density is zero. So the Hubble horizon is infinite at these turning points. We also find that the cosmic spacetime is geodesically complete for both time-like and null geodesics. Therefore, there is no Big-bang or Big-crunch singularity in the space-time. We calculate the particle horizon and the event horizon, respectively, and find they are infinite. So the horizon problem is vanishing. Using the Wheeler-DeWitt equation, we show the system is quantum stable although a negative scalar potential is present. Finally, we prove that the flatness of spacetime and the primordial scale-invariant power spectrum could be generated in this cyclic model which is consistent with the findings of Gratton et al [31] and Boyle et al. [32].

Acknowledgments

We thank the referee for the perceptive and insightful report which has improved the paper significantly. This

work is supported by the National Science Foundation of China under the Grant No. 10973014, No. 11373031 and the 973 Project (No. 2010CB833004).

-
- [1] R. H. Brandenberger, [arXiv:hep-th/1204.6108].
 - [2] A. Borde, A. H. Guth and A. Vilenkin, Phys. Rev. Lett. 90, 151301 (2003) [gr-qc/0110012].
 - [3] P. J. Steinhardt and N. Turok, Science 296, 1436 (2002); P. J. Steinhardt and N. Turok, Phys. Rev. D 65, 126003 (2002).
 - [4] L. Baum, P. H. Frampton, Phys. Rev. Lett. 98, 071301 (2007).
 - [5] R. H. Brandenberger, Phys. Rev. D 80, 043516 (2009); Y. Cai, D. A. Easson, R. H. Brandenberger, JCAP, 08(2012)020; C. Lin, R. H. Brandenberger, L. P. Levasseur, JCAP 1104:019,2011; T. Biswas, E. Gerwick, T. Koivisto, A. Mazumdar, Phys. Rev. Lett. 108, 031101 (2012); T. Biswas, A. S. Koshelev, A. Mazumdar, S. Y. Vernov, JCAP 08 (2012) 024; T. Biswas, E. Gerwick, T. Koivisto, A. Mazumdar, JCAP 1011:008,2010; T. Biswas, A. Mazumdar, W. Siegel, JCAP 0603 (2006) 009;
 - [6] G.F.R. Ellis and R. Maartens, Class. Quantum Grav. 21, 223 (2004) [gr-qc/0211082v4];
 - [7] J. D. Barrow, G. F. R. Ellis, R. Maartens, and C.G. Tsagas, Class. Quant. Grav. 20, L155 (2003) [gr-qc/0302094];
 - [8] S. del Campo, E. Guendelman, A.B. Kaganovich, R. Herrera and P. Labrana, Phys. Lett. B 699, 211 (2011), [arXiv:1105.0651];
 - [9] P. Wu and H. Yu, Phys. Rev. D 81, 103522 (2010) [arXiv:0909.2821];
 - [10] P.W. Graham, B. Horn, S. Kachru, S. Rajendran, and G. Torroba, [arXiv:1109.0282];
 - [11] M. P. Dabrowski, Ann. Phys. 248, 199 (1996) [gr-qc/9503017].
 - [12] R. R. Caldwell, Phys. Lett. B 545:23-29, 2002; R. R. Caldwell, Marc Kamionkowski and N. N. Weinberg, Phys. Rev. Lett. 91 (2003) 071301; H. Wei, R. G. Cai and D. F. Zeng, Class. Quant. Grav. 22, 3189 (2005) [hep-th/0501160]; H. Wei and R. G. Cai, Phys. Rev. D 72, 123507 (2005) [astro-ph/0509328].
 - [13] P. J. E. Peebles and B. Ratra, Ap. J. Lett. 325, L17 (1988); B. Ratra and P. J. E. Peebles, Phys. Rev. D 37, 3406 (1988); J. Frieman, C. Hill, A. Stebbins, I. Waga, Phys. Rev. Lett. 75 2077 (1995); R. R. Caldwell, R. Dave and P. J. Steinhardt, Phys. Rev. Lett. 80, 1582 (1998); P. G. Ferreira and M. Joyce, Phys. Rev. Lett. 79, 4740 (1997); see also C. Wetterich, Nucl. Phys. B 302, 668 (1988) and E. J. Copeland, A.R. Liddle, and D. Wands, Phys. Rev. D 57, 4686 (1998); X. Zhang, Phys. Lett. B 611 (2005) 1-7; X. Zhang, Mod. Phys. Lett. A 20 (2005) 2575;
 - [14] D. N. Spergel et al., Astrophys. J. Suppl. Ser. 170, 377 (2007); Morad Amarzguoui, O. Elgaroy, David F. Mota and T. Multamaki, Astrophys. J. 454, 707 (2006).
 - [15] S. Chandrasekhar, The Mathematical Theory of Black Holes, (Oxford University Press, New York, 1983).
 - [16] S. A. Hayward, S. Mukohyama, and M.C. Ashworth, Phys. Lett. A 256, 347 (1999); S.A. Hayward, Class. Quantum Grav. 15, 3147 (1998).
 - [17] D. Bak and S.J. Rey, Class. Quantum Grav. 17, L83 (2000).
 - [18] R.G. Cai and S.P. Kim, J. High Energy Phys. 02, 050 (2005).
 - [19] Y. Gong and A. Wang, Phys. Rev. Lett. 99, 211301 (2007).
 - [20] M. Akbar and R.G. Cai, Phys. Rev. D 75, 084003 (2007).
 - [21] R.G. Cai and L.M. Cao, Phys. Rev. D 75, 064008 (2007).
 - [22] R.G. Cai and L.M. Cao, Nucl. Phys. B 785, 135 (2007).
 - [23] M. Akbar and R.G. Cai, Phys. Lett. B 648, 243 (2007).
 - [24] A. Sheykhi, B. Wang and R.G. Cai, Phys. Rev. D 76, 023515 (2007).
 - [25] A. Sheykhi, B. Wang and R.G. Cai, Nucl. Phys. B 779, 1 (2007).
 - [26] R. C. Tolam, Phys. Rev. 37, 1639 (1931); R. C. Tolam, Phys. Rev. 38, 1758 (1931).
 - [27] E. Verlinde, arXiv:hep-th/0008140 ; R. G. Cai and Y. S. Myung, Phys. Rev. D 67, 124021 (2003) [arXiv:hep-th/0210272]; S. Nojiri, S. D. Odintsov and S. Ogushi, Int. J. Mod. Phys. A 17, 4809 (2002) [arXiv:hep-th/0205187]; T. Padmanabhan, Class. Quant. Grav. 19, 5387 (2002), gr-qc/0204019; Phys. Rept. 406, 49 (2005), gr-qc/0311036; A. V. Frolov and L. Kofman, JCAP 0305, 009 (2003) [arXiv:hep-th/0212327]; U. H. Danielsson, Phys. Rev. D 71, 023516 (2005) [arXiv:hep-th/0411172]; R. Bousso, Phys. Rev. D 71, 064024 (2005) [arXiv:hep-th/0412197]; M. Akbar and R.G. Cai, Phys. Rev. D 75, 084003 (2007).
 - [28] E. Babichev, V. Dokuchaev, Yu. Eroshenko, Phys. Rev. Lett. 93, 021102(2004).
 - [29] G. Izquierdo, D. Pavon, Phys. Lett. B 633, 420 (2006).
 - [30] B.S. DeWitt, Phys. Rev. 160, 1113 (1967); A. Vilenkin, Phys. Rev. D 50, 2581 (1994); C. Kiefer and B. Sandhofer, arXiv:0804.0672 [gr-qc]; J. J. Halliwell, in Proceedings of the 1990 Jerusalem Winter School on Quantum Cosmology and Baby Universes, ed. by S. Coleman, J.B. Hartle, T. Piran and S. Weinberg (World Scientific, Singapore, 1991).
 - [31] S. Gratton, J. Khoury, P. J. Steinhardt and N. Turok, Phys. Rev. D 69, 103505 (2004).
 - [32] L. A. Boyle, P. J. Steinhardt and N. Turok, Phys. Rev. Lett. 96, 111301 (2006).
 - [33] J. K. Erickson, D. H. Wesley, P. J. Steinhardt and N. Turok, Rev. D 69, 063514 (2004).
 - [34] P. J. Steinhardt and N. Turok, New. Astron. Rev. 49, 43 (2005).
 - [35] M. Li, Phys. Lett. B 724, 192(2013); T. Qiu, X. Gao, E. N. Sadridakis, Phys. Rev. D 88, 043525 (2013); A. Fertig et al., Phys. Rev. D 89, 103537 (2014); A. Ijjas et al., Phys. Rev. D 89(2014)123520.
 - [36] J. Garriga and V. F. Mukhanov, Phys. Lett. B 458 (1999) 219-225.
 - [37] D. Wands, Phys. Rev. D 60, 023507 (1999).
 - [38] F. Finelli and R. Brandenberger, Phys. Rev. D 65, 103522 (2002).